

DASYCLADALEAN GREEN ALGAE AND ASSOCIATED FORAMINIFERS IN MIDDLE TRIASSIC (LOWER AND MIDDLE MUSCHELKALK) CARBONATES OF THE SOUTH-EASTERN GERMANIC BASIN (UPPER SILESIA, POLAND)

Ioan I. BUCUR¹ & Michał MATYSIK^{2,3*}

¹Babeș-Bolyai University, Department of Geology and Center for Integrated Geological Studies,
Kogălniceanu 1, 400084 Cluj-Napoca, Romania; e-mail: ioan.bucur@ubbcluj.ro

²Michał Matysik Geoconsulting, Malachitowa 5/3, 30-798 Kraków, Poland

³Institute of Geological Sciences, Jagiellonian University,
Gronostajowa 3a, 30-387 Kraków, Poland;
e-mail: michał.matysik@uj.edu.pl

*Corresponding author

Bucur, I. I. & Matysik, M., 2020. Dasycladalean green algae and associated foraminifers in Middle Triassic (Lower and Middle Muschelkalk) carbonates of the south-eastern Germanic Basin (Upper Silesia, Poland). *Annales Societatis Geologorum Poloniae*, 90: 391–407.

Abstract: Middle Triassic dasycladalean algae occur in limestones and dolstones of the Diplopora Beds in Upper Silesia, a region representing the southern part of the Germanic Basin. The dasycladalean assemblage of this area was studied by Kotański at the end of the last century, mostly from dolomitized and weathered material. The relatively well-preserved specimens the authors found in thin sections from undolomitized strata allowed the description and revised taxonomical assignment of the algae. The identified microflora includes: *Diplopora annulatissima* Pia, 1920, *Diplopora annulata* (Schafhäutl, 1853), *Salpingoporella* cf. *krupkaensis* Kotański, 2013, *Physoporella prisca* Pia, 1912, *Physoporella* cf. *pauciforata* (Gümbel, 1872), *Holosporella?* sp. 1, and *Holosporella?* sp. 2. *Diplopora annulatissima* and *Favoporella annulata* Sokać, 1986 are regarded here as two different species. *Oligoporella elegans* Assmann ex Pia, 1931, extensively illustrated and described by Kotański (2013), is considered a junior synonym of *Oligoporella prisca* Pia, 1912, and transferred to the genus *Physoporella* emended by Grgasovic, 1995. Also, some of the dasycladalean algae (*Salpingoporella* cf. *krupkaensis*, *Holosporella?* sp. 1, and *Holosporella?* sp. 2) described here could represent new species, but more well-preserved material is necessary. The algae are accompanied by foraminiferal microfauna, which collectively indicate a late Anisian (Illyrian) age of the strata studied.

Key words: Dasycladaleans, green algae, benthic foraminifera, Middle Triassic, Muschelkalk, Poland.

Manuscript received 4 August 2020, accepted 17 October 2020

INTRODUCTION

Since the nineteenth century, when the first studies on Triassic calcareous algae were published, especially from the Alpine region (e.g., Schafhäutl, 1953; Gümbel, 1872, 1874; Benecke, 1876; Salomon, 1895), descriptions have been provided also of the “Nulliporae” (i.e., dasycladaleans) from the Upper Silesian region in the south-eastern Germanic Basin (e.g., Eck, 1862; Gümbel, 1872; Ahlburg, 1906). From the very beginning, the authors emphasized the special kind of preservation of the dasycladaleans from the Upper Silesian region, affected by complete dissolution

of the originally aragonitic skeleton, leaving only internal moulds, which are now filled with cement. More recently, papers on the Upper Silesian dasycladaleans were published by Pastwa-Leszczyńska and Śliwiński (1960), Kotański (1979, 1981, 1986, 1994), and Myszkowska (1992). An important monograph dedicated to these algae was published posthumously by Kotański (2013). In this paper, Kotański described and illustrated 21 dasycladalean species, including three new species. The largest part of the material from the above-mentioned studies in Upper Silesia is represented

by internal moulds and collected from dolomitized strata of the so-called “Diplopora Dolomite” or “Diplopora Beds”.

In the present paper, the authors describe seven dasycladalean species identified in thin sections from the Diplopora Beds in Upper Silesia, mainly from localities that have never been studied for algae and where the strata are undolomitized. Most of these algae were identified also by Kotański (2013) as weathered material. The aim of the present authors was to study and illustrate these algae in relatively well-preserved specimens and to revise their systematic assignment according to some recent studies (e.g., Grgasović, 1995; Grgasović and Sokač, 2003).

GENERAL SETTING

Palaeogeography

In Middle Triassic times, the semi-closed and tropical Germanic Basin was connected with the southerly located Tethys Ocean through three narrow, submeridional sea-ways, called the East Carpathian Gate, Silesian-Moravian Gate, and Western Gate (Fig. 1). The gates developed at the tips of the Tethyan rift system and were opening and closing diachronously from the east to west, as the Tethyan spreading center migrated westwards through time (Szulc, 2000). The Upper Silesia study region is located at the northern outlet of the Silesian-Moravian Gate and was bounded to the west by the Bohemian Massif, to the east by the Małopolska Massif, and to the north-east by an archipelago of several cliff-edged Devonian islands.

The sedimentary record in Upper Silesia was controlled in the first place by long-term tectonic evolution of the neighboring gate. The gate opened in the Olenekian, reached its maximal width in the Pelsonian (mid-Anisian), and closed with the Fassanian (early Ladinian). Consequently, the Lower Muschelkalk deposited during

the gradual gate opening is ~100 m thick and dominated by open-marine limestones, while the Middle and Upper Muschelkalk formed during progressive tectonic uplift of the area is half the thickness and more dolomitic-siliciclastic (Szulc, 2000; Matysik, 2016). Within this 7-Myr trend, the temporal facies development reflected superimposed third- to fifth-order relative sea-level oscillations, which created an alternation of transgressive and highstand systems tracts (Szulc, 2000; Matysik, 2014, 2016) with internal high-frequency depositional cycles (Matysik, 2019). The spatial facies distribution in each time interval was dictated by the morphology of the Upper Silesian carbonate platform, which dipped to the south-west and thus had shallower and sometimes more restricted facies in the eastern (Kraków-Silesia) area, compared to the western (Opole) district (Wyczółkowski, 1971; Matysik, 2014; Matysik and Surmik, 2016). Local geomorphic lows and highs (Matysik, 2016) as well as synsedimentary block tectonics (Szulc, 1989, 1993; Matysik and Szulc, 2019) modified this simple facies pattern, sometimes producing a mosaic record.

Stratigraphy

The Upper Silesian Muschelkalk is subdivided into nine lithostratigraphic formations (informally called ‘beds’; Assmann, 1944; Śliwiński, 1961) that more-or-less correspond to the systems tracts of four third-order transgressive-regressive, disconformity-bounded sequences (Fig. 2; Szulc, 2000). The first Muschelkalk sequence (15 m thick) is largely composed of transgressive ramp tempestites and fair-weather lime muds, directly capped by lagoonal to supratidal dolomitic facies (Lower Gogolin Beds). The second Muschelkalk sequence (40 m thick) starts with a very similar transgressive ramp succession (Upper Gogolin Beds and the lower half of the Olkusz Beds), which evolved with time into a nearly flat carbonate bank with peloid-oncoid-oolithic sand shoal complexes and interbedded micrites representing highstand deposits (Górażdże Beds and the upper half of the Olkusz Beds).

The third Muschelkalk sequence (50 m thick) was deposited during the maximum opening of the Silesian-Moravian Gate and is the main storage setting for green algae in the Upper Silesian Muschelkalk. The sequence displays a very complex internal architecture, with peritidal carbonates in the eastern Kraków-Silesia region and open-marine facies in the western Opole region (Fig. 2). The peritidal succession representing a back-barrier area is tripartite. Its lower (transgressive) and upper (regressive) parts comprise lagoonal peloid sands and burrowed muds, intertidal cryptalgal laminites and loferites, and supratidal caliche levels, organized in various configurations into metre-scale cycles (Matysik, 2019). The middle part, corresponding to maximum flooding of the area, is entirely subtidal and dominated by peloid and oncoid sands (Fig. 3A). All these strata are dolomitized and are called “Diplopora Beds” because of ubiquitous diploporean green algae, which are concentrated particularly in the middle subtidal part of the succession (Fig. 3B). In contrast to the peritidal succession, the open-marine succession, developed in front of the barrier, starts with transgressive tempestites and deeper-platform




Fig. 1. Palaeogeographic position of the Upper Silesia study region (white rectangle) in the Middle Triassic. Modified from Szulc (2000).





Fig. 2. Generalized stratigraphic section for the Upper Silesian Muschelkalk, showing thickness, overall lithological character, provisional formation names, 3rd-order transgressive-regressive sequences, and algae-rich intervals (red thick outlines). The transition between the eastern Kraków-Silesia region and western Opole region is not recognized because of the lack of outcrops. Note that the algae-bearing interval is slightly older in the Kraków-Silesia region than in the Opole region, on the basis of the sequence stratigraphic correlation. Scheme modified from Matysik (2014), sequence stratigraphy framework after Szulc (2000), supplemented by Matysik (2014), lithostratigraphy after Assmann (1944) and Śliwiński (1961) with later changes by Szulc (2000) and Matysik (2014).



lime muds (*Terebratula* Beds). They are succeeded by a regressive suite of micritic firmgrounds, bioclastic tempestites, sponge-scleractinian patch reefs, and circumreefal crinoid-peloid sands (Karchowice Beds), shallowing to ooid-peloid sands with abundant green algae (*Diplopora* Beds; Fig. 3C–E; Matysik, 2010). The latter are generally dolomitized in their upper part, with only few localities from the lower part avoiding dolomitization, which were the main focus of this study (Fig. 3C–H). The barrier separating the peritidal and open-marine domains is unknown due to the lack of outcrops and cores.

The last Muschelkalk sequence is poorly exposed. It comprises restricted dolomitic (Tarnowice Beds), transgressive open-marine (Wilkowice Beds), and regressive siliciclastic (Boruszowice Beds) facies, deposited in a marginal sea (Szulc, 2000).

The sequence stratigraphy framework outlined above is correlated with the Alpine successions using magnetostratigraphy (Nawrocki and Szulc, 2000) and conodont, ammonoid, crinoid, and green algae biostratigraphy (Assmann, 1944; Zawidzka, 1975; Haggdorn and Głuchowski, 1993; Kotański, 1994; Kaim and Niedzwiedzki, 1999; Narkiewicz and Szulc, 2004). Details of the stratigraphy, sedimentology, environmental setting, and ichnology are discussed in numerous articles, particularly in Assmann (1944), Szulc (2000), Matysik (2014, 2016), and Stachacz and Matysik (2020).

MATERIALS AND METHODS

Fieldwork was conducted in 15 large, active and small, inactive quarries, scattered over an area of 120 x 50 km between Opole in the west and Chrzanów in the east (for the GPS coordinates of each outcrop, see the Appendix). Each outcrop was subjected to a search for fossil algae and associated bioclasts in the walls and loose blocks. The most interesting sections were also measured bed by bed. Around 50 samples were collected and slabbed for further palaeontological and facies analyses. Petrographic observations of 61 thin sections were focused on identifying fossils and characterizing their host sediment. All thin sections collected are housed in the Institute of Geological Sciences, Jagiellonian University in Kraków.

RESULTS

Environmental context for the algae and their preservation

The algae studied generally occur in grainstones, where they are the main rock-forming component or the accessory component of peloids, ooids, oncoids, and/or other bioclasts, including bivalves, gastropods, crinoids, small benthic foraminifers, and ostracodes (Fig. 3). However, depositional and diagenetic aspects of the sediments vary across Upper Silesia, affecting the preservation of the algae.

In the eastern Kraków-Silesia region (peritidal back-barrier facies), the sediments are densely packed and partly to completely dolomitized (Matysik, 2014), which hinders the detailed taxonomical identification of the algae (Fig. 3B). Only rare specimens are better preserved. This kind of material prevails in the articles published so far (Pastwa-Leszczyńska and Śliwiński, 1960; Kotański, 1979, 1981, 1986, 1994, 2013; Myszkowska, 1992). The localities studied include Nowa Wioska, Ząbkowice, Libiąż, Stare Gliny, and Przelajka.

In contrast, in the western Opole region (open-marine facies), the sediments are less densely packed and often volumetrically dominated by various (mainly early marine) aragonite and calcite cements, which most likely inhibited the downward percolation of dolomitizing fluids from the overlying Tarnowice Beds. In most localities, including Tarnów Opolski, Kamień Śląski, and Strzelce Opolskie, the grains are coated with a <0.2-mm-thick isopachous rim of fibrous cement, with the remaining interparticle pore space filled by drusy calcite or calcite spar (Fig. 3E). Farska Kolonia is unique, because the components are often very loosely organized and enclosed by an isopachous rim of radiaxial-fibrous cement, 0.2–2 mm thick, filling most interparticle pores. In scattered larger pores (which could be growth-framework cavities), this cement is locally post-dated by botryoidal cement, with the remaining pore space occluded with drusy calcite (Fig. 3F–H). The loose character of sediments, lack of interparticle micrite, and variety of early-marine cements collectively point to a turbulent, mud-free habitat for the algae and very rapid cementation of the sediment, almost at the seafloor.

Fig. 3. Selected sedimentological aspects of the algae-rich *Diplopora* Beds in the Kraków-Silesia (A, B) and Opole (C–H) regions. **A.** Representative outcrop view of dolomitized *Diplopora* Beds, with lower and upper peritidal facies separated by thick-bedded, lagoonal, algae-rich grainstones. Nowa Wioska. **B.** Detail of A, showing green algae (white arrows), gastropods (black arrows), and bivalves (blue arrow), embedded in peloid grainstone. **C.** Outcrop view of sponge bioherm (Karchowice Beds), overlain by undolomitized *Diplopora* Beds, shallowing from peloid grainstone, to ooid grainstone, to peloid grainstone with abundant green algae. Kamień Śląski. **D.** Bedding plane view of ooid-peloid grainstone with bivalves (blue arrows), crinoids (red arrows), and green alga fragment (white arrow). The sediment comes from Tarnów Opolski, but corresponds to the level of pure ooid grainstone shown in C. **E.** Plane-polarized view of C, showing undetermined green algae (yellow arrows), peloids (white arrows), and small benthic foraminifer (red arrows), all enclosed by thin isopachous rim of fibrous cement (blue arrows). The remaining interparticle pore space in the upper half of the photo is filled with drusy calcite (dc). **F.** Vertically oriented slab of oncoid (o) grainstone with scattered green algae (white arrows) and various cements filling interparticle pores (red arrows). Farska Kolonia. **G, H.** Plane- and cross-polarized views of larger interparticle pore, lined with radiaxial-fibrous cement (rfc), post-dated by botryoidal cement with fibres growing perpendicular to the photo (bc), and occluded with drusy calcite (dc). Farska Kolonia.

Systematics

Order DASYCLADES Pascher, 1931

Family DIPLOPOREAE (Pia, 1920) Deloffre, 1988
 Genus *Diploporella* Schafhäutl, 1863, emend. Güvenç, 1979
Diploporella annulatissima Pia, 1920
 Fig. 4A–I

- 1920 *Diploporella annulatissima* nov. spec. – Pia, p. 67, text-fig. 15, plate 4, figs 11–16.
- 2000 *Kantia monregalensis* Barette, 1919 – Granier and Grgasović, p. 85, with extended synonymy.
- 2013 *Diploporella annulatissima* Pia, 1920 – Kotański, p. 96, text-fig. 23, plate 26–30.

Remarks: *Diploporella annulatissima* is one of the most frequent dasycladalean species in the region studied; this was also noted by Kotański (2013, p. 97). It is characterized by a single whorl of laterals within a ring (Fig. 4A–G). The laterals have a trichophorous to acrophorous shape (Fig. 4H, I), except for the specimen in Fig. 4C, which shows a possible distal enlargement, characteristic of *Favoporella annulata* Sokač, 1968. The tangential sections (Fig. 4F, G) show four laterals in a tuft.

Dimensions (in mm):

D (external diameter of the skeleton) = 2.04–2.94 (mean = 2.43);
 d = (internal diameter of the skeleton) = 1.09–1.79 (mean = 1.41);
 $d/D = 0.53–0.61$ (mean = 0.58);
 l (length of the laterals) = 0.41–0.63 (mean = 0.53);
 h (distance between verticils) = 0.38–0.62 (mean = 0.49);
 Thickness of a ring = 0.22–0.38 (mean = 0.28).


The possible synonymy between *D. annulatissima* and *Kantia monregalensis* Barette, 1919 was questioned by Herak (1957). This author (Herak, 1957, p. 49) states: “A comparison of some published figures of *Kantia monregalensis* (Barette, 1919, fig. 6) with those of the type specimens of *Diploporella annulatissima* (Pia, 1920, plate 4, figs 12–14) leaves no doubts that they are the same species.” But further on at the same page, Herak (1957) writes: “If *Kantia monregalensis* had been validly published, the name of the species would now be *Diploporella monregalensis* (Barette) as the name *Kantia* has been rejected. This is impossible, however, because heterogenous specimens were figured under the name *Kantia monregalensis*, and the description does not correspond to specimens that are now regarded as true representatives of the species in question. The present authors must therefore retain the name *Diploporella annulatissima*”. The genus *Kantia*, rejected by Pia (1920),

was re-introduced by Güvenc (1979) together with emendation of the genus *Diploporella*. The species assigned to the genus *Kantia* are characterized by vesiculiform laterals, while the species assigned to the genus *Diploporella* have trichophorous laterals. A third category of species, ascribed by some authors to the genus *Diploporella* characterized by phloio-phorous laterals (e.g., *Diploporella proba* Pia, 1935), has been transferred to the genus *Pseudodiploporella* (Bucur and Enos, 2001). It is noteworthy that illustrations provided by Barette (1919) are schematic drawings. A serious comparison between *Kantia monregalensis* and *Diploporella annulatissima* based on these illustrations is actually impossible because the laterals’ shape is not visible in these schematic representations. Moreover, as Barretti assigned the alga to the genus *Kantia*, the authors have to suppose that the laterals were vesiculiform. In fact, Barette (1919, p. 280) noted: “I pori... sono ingrossati notevolmente nella parte esterna (The pores... are enlarged significantly on the outside...)”. Consequently, the present authors believe that the synonymy between *Diploporella annulatissima* and *Kantia monregalensis* (Granier and Grgasović, 2000) is not appropriate.

Another problem related to *Diploporella annulatissima* is its possible synonymy with *Favoporella annulata*, which was stated by Ott (1972) and subsequently accepted by Grgasović and Sokač (2003). These authors considered that *D. annulatissima* and *F. annulata* represent two forms of the same alga, but with a different degree of calcification. Bucur and Enos (2001, p. 324) rejected this synonymy on the basis of the different shape of the laterals: trichophorous in *D. annulatissima* and phloio-phorous in *F. annulata*. In comparison of the two algae, two problems arise:

1. It is not clear if the arrangement of the laterals in *Favoporella annulata* is of metaspondyle type. Pictures provided by Sokač (1968), including his figure 1 with the alga reconstruction, rather show that the laterals do not start from the same insertion point of the main stem and accordingly it is not a metaspondyle alga. Besides, Sokač (1968, p. 210) noted this feature in the diagnosis of the species *F. annulata*: “The branches with the widened bases depart separately from the main stem...”. On the contrary, *D. annulatissima* has a metaspondyle arrangement (see for instance Bucur and Enos, 2001, plate 3, fig. 13; Kotański, 2013, plate 27, fig. 6, plate 28, fig. 1). This characteristic would differentiate both algae. But if the arrangement of the laterals in *F. annulata* proves to be metaspondyle, and we accept that *D. annulatissima* and *F. annulata* represent two forms with a different degree of calcification of the same alga, than this alga must be assigned to the genus *Pseudodiploporella* (phloio-phorous laterals with metaspondyle arrangement, Bucur and Enos, 2001) in the new combination *Pseudodiploporella annulatissima*. To resolve this issue, the careful study of well-preserved

Fig. 4. *Diploporella annulatissima* Pia, 1920. All specimens from Farska Kolonia. **A–D.** Longitudinal oblique (A, C, D) and oblique (B) sections. Note in C the external enlargement of the laterals (*Favoporella*-type laterals). A – thin section FaK4/952, B – thin section FaK4/954, C – thin section FaK/959, D – thin section FaK/961. **E.** Longitudinal section. Thin section FaK5/960. **F, G.** Tangential sections. F – thin section FaK5/960, G – thin section FaK5/961. **H.** Oblique section showing the shape of the laterals and their metaspondyle arrangement (arrow). Thin section FaK5/964. **I.** Fragment of transverse section showing the shape of the laterals. Thin section FaK4/953.



- specimens of *Favoporella annulata* is necessary, to see if the alga is metaspondyle or not.
2. It is noteworthy that Kotański (2013) raised the issue of possible synonymy between *D. annulatissima* and *Gyroporella cylindrica* Gümbel, 1872, but he also emphasized the necessity to formally conserve the name *D. annulatissima*.

Diplopora annulata (Schafhäutl, 1853) Schafhäutl, 1863
Fig. 5A–C

- 1853 *Nullipora annulata* n. sp. – Schafhäutl, plate 6, fig. 1a–f.
- 1863 *Diplopora annulata* n. gen., n. comb. – Schafhäutl, p. 324, plate 65e, fig. 6a, b.
- 2000 *Diplopora annulata* (Schafhäutl, 1853) 1863 – Granier et Grgasović, p. 32. [With extended synonymy.]

Remarks: Rare specimens belonging to this species were found in thin sections with frequent *Diplopora annulatissima*.

Family TRIPLOPORELLACEAE (Pia, 1920)
Berger et Kaever, 1992

Genus *Salpingoporella* (Pia, 1918)
emend. Carras et al., 2006

Salpingoporella cf. *krupkaensis* Kotański, 2013
Fig. 5D–K

- 2013 *Salpingoporella krupkaensis* n. sp. – Kotański, p. 50, plate 1–4.

Remarks: *Salpingoporella krupkaensis* was described by Kotański (2013, p. 50) as having "...Branches (tubercles) weakly attached to the wall of the central cavity, rapidly extended distally and froming funnel-like tubercles (plate 1, fig. 7) merging with the adjacent rock, where they begin to touch, forming probably a rarely and poorly preserved cortex (plate 2, fig. 4) absent from Jurassic and Cretaceous species (Kamptner, 1958, p. 110)". Kotański (2013) also remarked that *S. krupkaensis* is the oldest species of the genus *Salpingoporella*. However, he considered *S. krupkaensis* as "...the only Triassic species of *Macroporella* sensu lato with consequently euspondyle whorls..." (Kotański, 2013, p. 51), leading thus to a confusion between *Macroporella* and *Salpingoporella*. He did not compare *S. krupkaensis* with other Triassic, Jurassic or Cretaceous *Salpingoporella*, but with *Kantia comedicana*, an alga with a metaspondyle arrangement of vesiculiform laterals. Two comments should be made in connection with these remarks of Kotański: 1) by its euspondyle arrangement of laterals, *Salpingoporella* differs essentially from *Macroporella*, the two genera belonging to two different families of dasycladalean algae: euspondyle Triploporellaceae and aspondyle Seletonellaceae, respectively; 2) in contrast to the statement of Kamptner (1958, p. 110), all species of *Salpingoporella* are supposed to have an external cortex (see emended diagnosis of the genus *Salpingoporella* in Carras et al., 2006, p. 458).

Kotański (2013) gives no dimensions for his new species, considering it an unimportant feature. However, dimensions could be considered for discriminating species in some cases (cf. Bassoulet et al., 1977, p. 159). The authors measured the dimensions of the specimens illustrated by Kotański (2013), using the magnification in his photos. As emphasized in Table 1, the specimens from the material of the authors are smaller, hence the incertitude regarding the identity of the two algae. Six species of *Salpingoporella* are known from Upper Triassic deposits worldwide (cf. Schlagintweit et al., 2001; Carras et al., 2006): *Salpingoporella spectabilis* (Bystrický, 1962), *S. humilis* (Bystrický, 1967) *S. sturi* (Bystrický, 1967), *S. tibetica* Flügel and Mu, 1982, *S. triasica* Richter et al., 1999, and *S. austriaca* Schlagintweit et al., 2001. By its dimensions, *S. cf. krupkaensis* is closer to *S. triasica* Richter et al., but it differs from the latter by its larger mean values of the main parameters, by the much larger laterals (larger pores), and by the thicker calcareous wall. *S. austriaca* is even smaller and has closely set verticils of conical laterals. All other Triassic *Salpingoporella* differ from *S. cf. krupkaensis* by their overall larger dimensions.

On the basis of the dimensions, which do not overlap in any parameter, the present authors deduce that *S. krupkaensis* Kotański, 2013 and *S. cf. krupkaensis* are two different species. However, taking into account their close morphological similarity, the authors prefer to ascribe the identified specimens, with some reservations to *Salpingoporella krupkaensis*.

Table 1

Comparison between the general dimensions of *Salpingoporella krupkaensis* Kotański, 2013 and *Salpingoporella* cf. *krupkaensis* described in this paper

	<i>Salpingoporella krupkaensis</i> Kotański, 2013	<i>Salpingoporella</i> cf. <i>krupkaensis</i> (this paper)
L (max. obs.)	5.83	2.69
D	1.00–1.27	0.44–0.72 (mean = 0.58)
d	0.50–0.66	0.18–0.50 (mean = 0.29)
h	0.17–0.22	0.24
l	0.19–0.22	0.09–0.18 (mean = 0.14)
p (distal)	–	0.09–0.14 (mean = 0.12)
w	–	8–9

Genus *Physoporella* Steinmann, 1903 emend.
Grgasović, 1995
Physoporella prisca Pia, 1912
Fig. 6A–G

- non 1912 *Oligoporella prisca* mihi – Pia, p. 42, plate 5(4), figs 2, 4, 6, 8.
- 1926 *Diplopora elegans* n. sp. – Assmann, p. 505. [Nomen nudum, type not illustrated.]
- 1931 *Oligoporella elegans* Assmann – Pia, p. 273, plate 21, figs 3–6.




Fig. 5. Dasycladalean algae. A–C. *Diplopora annulata* (Schafhäutl, 1853) in longitudinal-oblique section (A), oblique-tangential section (B), and transverse section (C). Farska Kolonia. A – thin section FaK5/962, B, C – thin section FaK5/960. D–K. *Salpingoporella* cf. *krupkaensis* Kotański, 2013 in longitudinal-oblique section (D), oblique sections (E, F, H, K), longitudinal section (G), and transverse sections (I, J). D, E, G – Farska Kolonia, thin section FaK5/960, F, H, I, J – Tarnów Opolski, thin section TOp/oolity, K – Strzelce Opolskie, thin section SO1/101.

- 2000 *Oligoporella elegans* Assmann ex Pia 1931.
[= *Diploporella elegans* Assmann 1926, nomen nudum – Granier et Grgasović, p. 111, with synonymy.]
- 2000 *Oligoporella prisca* Pia 1912 – Granier et Grgasović, p. 119. [with synonymy.]
- 2013 *Oligoporella elegans* Assmann ex Pia 1931a – Kotański, p. 52, text-fig. 2, plates 5–7.

Remarks: Validating the species *Oligoporella elegans* Assmann 1926, Pia (1931, p. 273) noted (translated from German): “*Oligoporella elegans* is very close to *O. prisca*. The Alpine species has larger mean dimensions, so that the smaller specimens reach the dimensions of the German specimens. The internal cavity is smaller in *Oligoporella prisca* (with respect to D) and the verticils are closer. In general, they can be very numerous. The occasional enlargement of the pores towards the exterior is present in both species. It is better, however, to keep both species separate, at least until localities with mixed specimens will be found”. In its revision of the genera *Physoporella* and *Oligoporella*, Grgasović (1995) synonymized the species *O. elegans* and *O. prisca*, with the latter species as senior synonym being assigned to the emended genus *Physoporella*.

Kotański (2013) gives the following dimensions of *Physoporella prisca* (sub *Oligoporella elegans*): D = 0.58–1.0 mm; d = 0.27–0.64 mm; D/d = 46–63%; h = 0.2 mm. The dimensions we measured on the specimens studied in this paper are:

$$\begin{aligned} D &= 0.34–0.90 \text{ (mean = 0.65) mm;} \\ d &= 0.17–0.54 \text{ (mean = 0.37) mm;} \\ D/d &= 46.6–66.6\%; \\ h &= 0.17–0.45 \text{ (mean = 0.31) mm.} \end{aligned}$$

Physoporella cf. pauciforata (Gümbel, 1972)

Steinmann, 1903

Fig. 6H–O

- 1872 *Gyroporella pauciforata* n. sp. – Gümbel, p. 274, plate D3, fig. 2a–e.
- 1903 *Physoporella pauciforata* (Gümbel) nov. comb. – Steinmann, p. 50.
- 2000 *Physoporella pauciforata* (Gümbel 1872) Steinmann, 1903 – Granier et Grgasović, p. 130 (with extended synonymy).

Remarks: Together with *Physoporella prisca* we found specimens which are morphologically similar to *Physoporella pauciforata*, but of smaller dimensions:

D = 0.84–1.27 (mean = 1.02) mm;
d = 0.40–0.91 (mean = 0.61) mm;
h = 0.28–0.32 (mean = 0.29) mm.

Physoporella pauciforata typically has larger dimensions (e.g., Gümbel, 1872; Bystrický, 1964; Schlagintweit *et al.*, 2003).


Genus *Holosporella* Pia, 1930

Holosporella? sp. 1

Figs 7A–C, 8A, B

Remarks: A single thin section from Farska Kolonia provided numerous specimens of spheroidal-ovoidal bodies, sometimes presenting a small peduncle. They have a thin, micritic, unperforated wall. Elliott (1958) described from the Upper Triassic of Iraq such spherical bodies as a new problematic fossil, *Lacrymorphus perplexus*, and emphasized that some of the specimens have a pear, acorn or retort shape. Further species of *Lacrymorphus* have been described from Cretaceous strata by Radoičić (1967) as *Lacrymorphus catenaformis* and *Lacrymorphus globosus*. The first one was subsequently transferred to the genus *Humiella* Sokač and Velić, 1981 (Masse *et al.*, 1984). Similar microfossils were illustrated by Pecorini (1972) as *Lacrimorphus catenaformis* Radoičić *sardus* n. subsp. Cherchi and Schroeder (1985) revised these fossils and assigned them to the genus *Sarfatiella* Conrad and Peybernès, 1974 under the combination *Sarfatiella sarda*. Subsequently, *Sarfatiella* became a junior synonym of the genus *Holosporella* Pia, 1930. Following Masse *et al.* (1984), the two genera are differentiated by the presence (in *Humiella*) or absence (in *Sarfatiella* [= *Holosporella*]) of perforations in the wall of ampoule-like laterals. It is worth mentioning that pores in the calcareous coating of dasycladaleans also have been noted in *Kopetdagaria sphaerica* Maslov by Bucur (2002) and in *Actinoporella podolica* by Granier (1988) and Schlagintweit and Enos (2012). Following Schlagintweit and Enos (2012), this should be a species-dependent character. Starting from these considerations, the spherical-ovoidal bodies from Upper Silesia could belong to laterals of *Holosporella*. Together with the spherical bodies we found some rare dasycladalean skeletons that preserve a calcareous wall around the central cavity, which appears to be the proximal part of laterals (Fig. 7C). It is possible that these algae bore the spherical bodies (fertile ampoules) like in the hypothetical reconstruction in Figure 8A. Another possibility is that no relationship existed between these two algae and the spherical bodies simply represent fertile ampoules of *Holosporella*, as in the hypothetical reconstruction in Figure 8B. *Holosporella?* sp. 1 differs from the Upper Triassic *Lacrymorphus perplexus* by its much larger dimensions.

Fig. 6. Dasycladalean algae. A–G. *Physoporella prisca* Pia, 1912. A. Longitudinal and transverse oblique sections. Thin section FaK4/954. B. Close-up view of the specimen in longitudinal section in A. C. Longitudinal and transverse sections. Thin section FaK4/952. D. Longitudinal-oblique section. Thin section FaK5/959. E. Transverse section. Thin section FaK5/960. F. Close-up view of the specimen down-left in A. G. Oblique section. Thin section FaK5/960. All specimens from Farska Kolonia. H–O. *Physoporella cf. pauciforata* (Gümbel, 1872) in oblique sections. All specimens from Farska Kolonia. H, L – thin section FaK4/962, I – thin section FaK4/950, J, N, O – thin section FaK4/952, K – thin section FaK4/951, M – thin section FaK5/960.






Fig. 7. Dasycladalean algae. **A–C.** *Holosporella?* sp. 1. A – spheroidal and ovoidal bodies spread in the sediment, probably representing fertile ampoules of a *Holosporella*. Thin section FaK4/949. B – close-up view of the specimen in the middle of A, showing a short peduncle. C – Dasycladalean alga associated with the spheroidal-ovoidal bodies. Thin section FaK4/949. All specimens from Farska Kolonia. **D–F.** *Holosporella?* sp. 2. in oblique section (D), longitudinal section (E), and transverse section (F). All specimens from Farska Kolonia, thin section FaK4/950.

Holosporella? sp. 2
Fig. 7D–F

Remarks: In two thin sections from Farska Kolonia, there were rare specimens of a dasycladalean alga, which probably had vesiculiform laterals. Calcification affected only the proximal part of the laterals, so that the distal part cannot be observed. Somewhat similar specimens were recently described from the Upper Triassic of North America as *Holosporella?* *rossanae* Bucur and Del Pierro (in Bucur *et al.*, 2020), but the Upper Silesian specimens have slightly larger dimensions:

$$\begin{aligned} D &= 0.86–1.14 \text{ (mean = 1.02) mm;} \\ d &= 0.50–0.64 \text{ (mean = 0.59) mm;} \\ p &= 0.19–0.24 \text{ (mean = 0.22) mm;} \\ h &= 0.31 \text{ mm.} \end{aligned}$$

Only three species of *Holosporella* are known from Triassic deposits: *Holosporella conradii* Barattolo *et al.*, 2008, *Holosporella magna* Bucur and Fucelli (in Bucur *et al.*, 2020), and *Holosporella?* *rossanae* Bucur and Del Pierro (in Bucur *et al.*, 2020). As mentioned before,

Holosporella? sp. 2 from Silesia is closer to *H.?* *rossanae*, but more material is necessary to compare the two species properly.

Associated microfossils and age of the alga-bearing deposits

The calcareous algae are associated with the following benthic foraminifera (Fig. 9): *Endotriadiella wirzi* (Koehn-Zaninetti, 1969) (Fig. 9A, I, J), *Earlandinita elongata* Salaj in Salaj *et al.*, 1967 (Fig. 9B), *Endotebanella* sp. (Fig. 9C, D), *Meandrospira deformata* Salaj, 1967 (Fig. 9E), *Meandrospira dinarica* Kochansky-Devidé and Pantić (Fig. 9F–H), *Pilammina densa* Pantić (Fig. 9K), *Trochammina cf. alpina* Kristan-Tollmann, 1964 (Fig. 9L), *Tolyphammina* sp. (Fig. 9M), and *Diplotrema astrotuberculata* Kristan-Tollmann, 1960 (Fig. 9N). The foraminiferal assemblage, along with frequent *Diplopora annulatissima*, indicates a late Anisian (Illyrian) age of the Diplopora Beds (Salaj *et al.*, 1983, 1988).




Fig. 8. Two hypothetical reconstructions of *Holosporella* sp. 1. **A.** Reconstruction based on the supposed relationship between the spherical bodies and the associated dasycladalean. The reconstruction is a photomontage, using the dasycladalean in Figure 7C and spherical body in Figure 7B. **B.** Alternative reconstruction showing the arrangement of the fertile ampoules around a non-calcified main stem. The reconstruction is a photomontage, using the spherical body in Figure 7B. **C.** Schematic drawing of A. **D.** Schematic drawing of B.

CONCLUSIONS

The Middle Triassic carbonate deposits of the south-eastern Germanic Basin (Upper Silesia) contain in their upper part (so-called Diplopora Beds) numerous specimens of dasycladalean algae. The algae were studied extensively by Kotański (2013) on weathered material. The supplementary study by the present authors on relatively well-preserved specimens from undolomitized strata revealed seven species and allowed the revision of their systematic position. The present authors consider *Diplopora anulatissima* Pia, 1920 and *Favoporella annulata* Sokač, 1968 as two different species, at least until the metaspondyle arrangement of the laterals in *Favoporella* will be proven. *Oligoporella elegans* Assmann ex Pia, 1931 is considered a junior

synonym of *Oligoporella prisca* Pia, 1912 and is transferred to the genus *Physoporella* Steinmann, 1903, emend. Grgasović, 1995. Some of the dasycladalean algae described here (*Salpingoporella* cf. *krupkaensis*, *Holosporella*? sp. 1, and *Holosporella*? sp. 2) could represent new species, but more well-preserved material is necessary to decide in this respect.

ACKNOWLEDGEMENTS

The material analysed comes from the Ph.D. dissertation of M. Matysik, which was funded by the Ministry of Science and Higher Education (Research Grant N307 119938) and supervised by the late Joachim Szulc, who always supported him with inspiring discussions and constantly motivated him to consider




Fig. 9. Foraminifera associated with dasycladalean algae. **A, I, J.** *Endotriadella wirzi* Koehn-Zaninetti, 1969. Tarnów Opolski, thin section TOp/576–577. **B.** *Earlandinita elongata* Salaj, 1967. Farska Kolonia, thin section FaK4/952. **C, D.** *Endotebanella* sp. C – Strzelce Opolskie, thin section SO4/326; D – Tarnów Opolski, thin section Top10/alg. **E.** *Meandrospira deformata* Salaj, 1967. Farska Kolonia, thin section FaK4/953. **F–H.** *Meandrospira dinarica* Kochanski-Devidé & Pantić, 1966. Libiąż, thin section Li8. **K.** *Pilammina densa* Pantić, 1965. Farska Kolonia, thin section FaK4/953. **L.** *Trochammina* cf. *alpina* Kristan-Tollmann, 1964. Farska Kolonia, thin section FaK4/956. **M.** *Tolypammina* sp. Farska Kolonia, thin section FaK4/952. **N.** *Diplotremmina astrofimbriata* Kristan-Tollmann, 1960. Farska Kolonia, thin section FaK4/954.

a subject in depth. Reviewers Felix Schlagintweit and Jochen Kuss are thanked for critical and valuable comments on the manuscript.

REFERENCES

- Ahlburg, J., 1906. Trias im südlichen Oberschlesien. *Ahandlungen der Königlich Preußischen Geologischen Landesanstalt und Bergakademie, Neue Folge*, 50: 1–163.
- Assmann, P., 1926. Die Fauna der Wirbellosen und Diploporen der oberschlesischen Trias mit Ausnahme der Brachiopoden, Lamellibranchiaten, Gastropoden und Korallen. *Jahrbuch der Preußischen Geologischen Landesanstalt zu Berlin für das Jahr 1925*, 46: 504–527.
- Assmann, P., 1944. Die Stratigraphie der oberschlesischen Trias. *Teil II – Der Muschelkalk. Abhandlungen des Reichsbergs für Bodenforschung*, 208: 1–124.
- Baretti, A., 1919. Contributo allo studio delle Siphoneae Verticillatae del Calcare di Vilanova-Mondovi. *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale in Milano*, 58: 216–236.
- Bassoulet, J. P., Bernier, P., Deloffre, R., Génot, P., Jaffrezo, M., Poignant, A. F. & Segonzac, G., 1977. Classification criteria of fossil Dasycladales. In: Flügel, E. (ed.), *Fossil Algae*. Springer, Berlin-Heidelberg, pp. 154–166.
- Benecke, E. W., 1876. Über einige Versteinerungen, insbesondere aus den Umgebungen von Esino. In: Über die Umgebungen von Esino in der Lombardei. *Geognostisch-paläontologische Beiträge*, 2(3): 296–317.
- Berger, S. & Kaever, M. J., 1992. *Dasycladales. An illustrated monograph of a fascinating algal order*. Georg Thieme Verlag, Stuttgart, 247 pp.
- Bucur, I. I., 2002. New remarks on *Kopetdagaria sphaerica* Maslov, 1960. In: Bucur, I. I. & Filipescu, S. (eds), *Research Advances in Calcareous Algae and Microbial Carbonates*. Cluj University Press, Cluj, pp. 83–94.
- Bucur, I. I. & Enos, P., 2001. Middle Triassic dasyclad algae from Guizhou, China. *Micropaleontology*, 47: 317–338.

- Bucur, I. I., Rigaud, S., Del Piero, N., Fucelli, A., Heerwagen, E., Peybernes, C., Peyrotte, G., Verard, C., Chablais, J. & Martini, R., 2020. Upper Triassic calcareous algae from the Panthalassa Ocean. *Rivista Italiana di Paleontologia e Stratigrafia (Research in Paleontology and Stratigraphy)*, 126: 499–540.
- Bystrický, J., 1962. New Dasycladaceae of the Triassic in the Slovak karst. *Geologický Sborník*, 13: 227–240.
- Bystrický, J., 1964. *Slovensky kras. Stratigrafia a Dasycladaceae mezozoika slovenského krasu*. Ustredný ústav geologický, Bratislava, 204 pp.
- Bystrický, J., 1967. Die obertriadischen Dasycladaceen der Westkarpaten. *Geologický Sborník*, 18: 285–309.
- Carras, N., Conrad, M. A. & Radoičić, R., 2006. *Salpingoporella*, a common genus of Mesozoic Dasycladales (calcareous green algae). *Revue de Paléobiologie*, 25: 457–517.
- Cherchi, A. & Schroeder, R., 1985. *Sarfatiella sarda* (Pecorini, 1972) n. comb (Dasycladaceae) from the basal Lower Cretaceous of NW Sardinia and some comments on *Lacymorphus* Radoičić and *Humiella* Sokač & Velić. *Bollettino della Società Paleontologica Italiana*, 23: 355–360.
- Conrad, M. A. & Peyerbèrs, B., 1974. Sur quelques Dasycladaceées (Chlorophycées) du Dogger des Pyrénées centrales et orientales franco-espagnoles. *Archives des Sciences Genève*, 26 (for 1973): 297–308.
- Deloffre, R., 1988. Nouvelle taxonomie des algues Dasycladales. *Bulletin des Centres de Recherche Exploration-Production Elf-Aquitaine*, 12: 165–217.
- Eck, H., 1862. Ueber den opatowitzer Kalkstein des oberschlesischen Muschelkalks. *Zeitschrift der deutschen geologischen Gesellschaft*, 14: 288–311.
- Elliott, G. F., 1958. Fossil microproblematica from the Middle East. *Micropaleontology*, 4: 419–428.
- Flügel, E. & Mu, X., 1982. Upper Triassic Dasycladaceae from Eastern Tibet. *Facies*, 6: 59–74.
- Granier, B. R. C., 1988. Algues Chlorophyceae du Jurassique terminal et du Crétacé inférieur en Alicante. *Mediterranea, Serie de Estudios Geológicos*, 5: 5–96.
- Granier, B. R. C. & Grgasović, T., 2000. Les algues dasycladales du Permien et du Trias. Nouvelle tentative d'inventaire bibliographique, géographique et stratigraphique. *Geologia Croatica*, 53: 1–197.
- Grgasović, T., 1995. Revision of genera *Physoporella* Steinmann and *Oligoporella* Pia (Dasycladales, green algae). *Sixth International Symposium on Fossil Algae and Carbonate Platforms*, Ankara, Abstracts, p. 15.
- Grgasović, T. & Sokač, B., 2003. Review of the fossil dasycladalean association in the Triassic of Croatia. *Acta Micropalaeontologica Sinica*, 20: 75–79.
- Gümbel, C. W., 1872. Die sogenannten Nuliporen (*Lithothamnium* und *Diplopora*) und ihre Beteiligung an der Zusammensetzung der Kalkgesteine. Zweiter Theil: Die Nulliporen des Thierreichs (*Dactyloporidae*) nebst Nachtrag zum ersten Theile. *Abhandlungen der mathematisch-physikalischen Klasse der königlichen bayerischen Akademie der Wissenschaften*, 11: 231–289.
- Gümbel, C. W., 1874. Ueber neue Gyroporellen aus dem Gailthaler Gebirge. *Verhandlungen der kaiserlich-königlichen geologischen Reichsanstalt*, 4: 79–80.
- Güvenç, T., 1979. Dasycladacees métaspordyles du Paléozoïque supérieur et du Trias. *Bulletin des Centres de Recherche Exploration-Production Elf-Aquitaine*, 3: 625–637.
- Hagdorn, H. & Głuchowski, E., 1993. Palaeobiogeography and stratigraphy of Muschelkalk echinoderms (Crinoidea, Echinoidea) in Upper Silesia. In: Hagdorn, H. & Seilacher, A. (eds), *Muschelkalk. Schöntaler Symposium 1991*. Korb, Stuttgart, pp. 165–176.
- Herak, M., 1957. Some problems in the dasyclad genus *Diplopora*. *Micropaleontology*, 3: 49–52.
- Kaim, A. & Niedźwiedzki, R., 1999. Middle Triassic ammonoids from Silesia, Poland. *Acta Palaeontologica Polonica*, 44: 93–115.
- Kamptner, E., 1958. Über das System und die Stammesgeschichte der Dasycladacean (Siphonae Verticillatae). *Annalen des Naturhistorischen Museums in Wien*, 62: 95–122.
- Kochansky-Devidé, V. & Pantić, S., 1966. *Meandrospira* in der unteren und mittleren Trias sowie einige begleitende Fossilien in den Dirariden. *Geološki vjesnik*, 19: 15–28.
- Koehn-Zaninetti, L., 1969. Les foraminifères du Trias de la région de l'Almtal (Haute Autriche). *Jahrbuch der geologischen Bundesanstalt, Sonderband*, 14: 1–155.
- Kotański, Z., 1979. Macroflora. Gromada Chlorophyta. Rodzina Dasycladaceae. In: Malinowska, L. (ed.), *Budowa geologiczna Polski. III. Atlas skałieniałości przewodniczących i charakterystycznych. 2a. Mezozoik. Trias*. Instytut Geologiczny, Wydawnictwa Geologiczne, Warszawa, pp. 215–222. [In Polish.]
- Kotański, Z., 1981. Nowe obserwacje nad Dasycladaceami z dolomitu diploporeowego z wierceń okolic Zawiercia (Wyżyna Śląsko-Krakowska). In: *Fauna i flora triasu obrzeżenia Górz Świętokrzyskich i Wyżyny Śląsko-Krakowskiej. Materiały V Krajowej Konferencji Paleontologów*. Kielce-Sosnowiec, pp. 74–78. [In Polish.]
- Kotański, Z., 1986. Macroflora. Class Chlorophyta. In: Malinowska, L. (ed.), *Geology of Poland. III. Atlas of Guide and Characteristic Fossils. Part 2a. Mesozoic. Triassic*. Instytut Geologiczny, Wydawnictwa Geologiczne, Warszawa, pp. 176–184.
- Kotański, Z., 1994. Middle Triassic Dasycladacea of the Upper Silesian – Cracow Region and their stratigraphical and palaeoecological significance. *Third International Meeting of Peri-Tethyan Epicratonic Basins. Cracow (Poland). Excursion Stuttgart, Guidebook*. Polish Geological Institute, Warszawa, pp. 59–66.
- Kotański, Z., 2013. Anisian Dasycladales from Upper Silesia and adjacent regions. In: Granier, B. & Peryt, T. (eds), *Carnets de Géologie, Book 2*, 206 pp.
- Kristan-Tollmann, E., 1960. Rotaliidea (Foraminifera) aus der Trias der Ostalpen. *Jahrbuch der geologischen Bundesanstalt, Sonderband*, 5: 47–78.
- Kristan-Tollmann, E., 1964. Foraminiferen aus den Rhätischen Zlambachmergeln der Fischerwiese bei Aussee im Salzkammergut. *Jahrbuch der geologischen Bundesanstalt, Sonderband*, 10: 1–189.
- Masse, J.-P., Aquaviva, M. & Luperto-Sinni, E., 1984. *Humiella catenaformis* (Radoičić), nov. comb. (Dasycladale) de l'Éocétacé italo-dinarique. *Revue de Micropaléontologie*, 27: 139–143.
- Matysik, M., 2010. Reefal environments and sedimentary processes of the Anisian Karchowice Beds in Upper Silesia,

- southern Poland. *Annales Societatis Geologorum Poloniae*, 80: 123–145.
- Matysik, M., 2014. Sedimentology of the “ore-bearing dolomite” of the Kraków-Silesia region (Middle Triassic, southern Poland). *Annales Societatis Geologorum Poloniae*, 84: 81–112.
- Matysik, M., 2016. Facies types and depositional environments of a morphologically diverse carbonate platform: a case study from the Muschelkalk (Middle Triassic) of Upper Silesia, southern Poland. *Annales Societatis Geologorum Poloniae*, 86: 119–164.
- Matysik, M., 2019. High-frequency depositional cycles in the Muschelkalk (Middle Triassic) of southern Poland: origin and implications for Germanic Basin astrochronological scales. *Sedimentary Geology*, 383: 159–180.
- Matysik, M. & Surmik, D., 2016. New occurrence of vertebrate remains within the Lower Muschelkalk (Anisian) peritidal deposits of the “Stare Gliny” quarry near Olkusz (Kraków-Silesia region, southern Poland). *Przegląd Geologiczny*, 64: 495–503. [In Polish, with English summary.]
- Matysik, M. & Szulc, J., 2019. Shallow-marine carbonate sedimentation in a tectonically mobile basin, the Muschelkalk (Middle Triassic) of Upper Silesia (southern Poland). *Marine and Petroleum Geology*, 107: 99–115.
- Myszkowska, J., 1992. Lithofacies and sedimentation of Diplopora Dolomite (Middle Muschelkalk) in the east part of Cracovian-Silesian region. *Annales Societatis Geologorum Poloniae*, 62: 19–62. [In Polish, with English summary.]
- Narkiewicz, K. & Szulc, J., 2004. Controls on migration of conodont fauna in peripheral oceanic areas. An example from the Middle Triassic of the Northern Peri-Tethys. *Geobios*, 37: 425–436.
- Nawrocki, J. & Szulc, J., 2000. The Middle Triassic magnetostratigraphy from the Peri-Tethys Basin in Poland. *Earth and Planetary Science Letters*, 182: 77–92.
- Ott, E., 1972. Neufunde anesischer Dasycladaceen (Kalkalgen) in Österreich. *Anzeiger der mathematisch-naturwissenschaftlichen Klasse der Österreichische Akademie der Wissenschaften*, 1972, 9: 1–14.
- Pantić, S., 1965. *Pilammina densa* n. gen., n. sp., and other Ammodiscidae from the Middle Triassic in the Crmnica (Montenegro). *Geološki vjesnik*, 18 (for 1964): 189–193.
- Pascher, A., 1931. Systematische Übersicht über die mit Flagellaten in Zusammenhang stehenden Algenreihen und Versuch einer Einreichung dieser Algenstämme in die Stämme des Pfanzenreiches. *Beihefte zum Botanischen Zentralblatt*, 48: 317–332.
- Pastwa-Leszczyńska, C. & Śliwiński, S., 1960. Algae (Dasycladaceae) from ore-bearing dolomites in the vicinity of Chrzanów. *Kwartalnik Geologiczny*, 4: 679–699. [In Polish, with English summary.]
- Pecorini, G., 1972. Microflora “purbekiana” della Nurra (Sardegna). *Bollettino della Società Geologica Italiana*, 91: 373–385.
- Pia, J., von, 1912. Neuen Studien über die triadischen Siphoneae Verticillatae. *Beiträge zur Paläontologie und Geologie Österreich-Ungarns und des Orients. Mitteilungen des Paläontologischen und Geologischen Institutes der Universität Wien*, 25: 25–81.
- Pia, J. von, 1918. Dasycladaceae. In: Trauth, F. (ed.), *Das Eozänvorkommen bei Radstadt im Pongau und seine Beziehungen zu den Gleichalterigen Ablagerungen bei Kirchberg am Wechsel und Wimpassing am Leithagebirge. Kaiserliche Akademie der Wissenschaften*, 95: 209–213.
- Pia, J. von, 1920. Die Siphoneae Verticillatae vom Karbon bis zur Kreide. *Abhandlungen der zoologisch-botanischen Gesellschaft in Wien*, 21: 1–263.
- Pia, J. von, 1930. Upper Triassic fossils from the Burmo-Siamese frontier – a new dasycladacea *Holosporella siamensis* nov. gen., nov. sp., with a description of the allied genus *Aciculella*. *Pia. Records of the Geological Survey of India*, 63: 177–181.
- Pia, J. von, 1931. Die Dasycladaceen der germanischen Trias. *Annalen des naturhistorischen Museums in Wien*, 45: 265–279.
- Pia, J., v., 1935. Die Diploporen der anischen Stufe Bosniens. *Geoloski Analji Balkanskog Poluostrva*, 12: 190–246.
- Radoičić, R., 1967. On problematic microfossils from the Jurassic and Cretaceous of the Dinarides. *Vesnik Zavoda za Geološka I geofizička Istraživanja. Serie A*, 24–25: 269–279.
- Richter, D. K., Dragastan, O., Kube, B., Beck, T. & Zuhl, H., 1999. Facies und Biostratigraphie des finalen Stadiums der obertriadischen Karbonatplatform von Hydra (Kap Kastello/Weststrand des Pelaginikums/Griechenland). *Bochumer Geologische und Geotechnische Arbeiten*, 53: 127–148.
- Salaj, J., Biely, A. & Bystrický, J., 1967. Trias-Foraminiferen in den Westkarpaten. *Geologické Práce, Zpravy*, 42: 119–136.
- Salaj, J., Borza, K. & Samuel, O., 1983. *Triassic foraminifers of the West Carpathians*. Geologicky Ustav Dionyza Stura, Bratislava, 213 pp.
- Salaj, J., Trifonova, E. & Gheorghian, D., 1988. A biostratigraphic zonation based on benthic foraminifera in the Triassic deposits of the Carpatho-Balkans. *Revue de Paléobiologie*, 2: 153–159.
- Salomon, W., 1895. Geologische und palaeontologische Studien über die Marmolata. *Palaeontographica. Beiträge zur Naturgeschichte der Vorzeit*, 42: 1–210.
- Schafhäutl, K. E., 1853. Beiträge zur Kenntnis der Bayerischen Voralpen. *Neues Jahrbuch für Mineralogie, Geologie, Geognosie und Petrafakten-Kunde. I. Abhandlungen*: 229–319.
- Schafhäutl, K. E., 1863. *Südbayerns Lethaea Geognostica. Der Kressenberg und südlich von ihm gelegenen Hochalpen geognostisch betrachtet in ihren Petrefacten*. L. Voss Verlag, Leipzig, 487 pp.
- Schlagintweit, F. & Enos, P., 2012. Uppermost Jurassic? – Neocomian shallow-water carbonates of the Black Nose, USA: DSDP Site 392A revisited. *Acta Palaeontologica Romaniae*, 9: 39–56.
- Schlagintweit, F., Mandl, G. W. & Ebli, O., 2001. *Salpingoporella austriaca* n. sp., a new dasycladale (calcareous alga) from the lower Norian Dachstein Limestone of Mount Dachstein (Northern Calcareous Alps, Austria). *Revue de Paléobiologie*, 20: 359–365.
- Schlagintweit, F., Rünstler, H. & Gawlick, H.-J., 2003. Über ein Vorkommen von *Physoporella pauciforata* (Gümbel 1872) Steinmann 1903 (Dasycladaceae) aus Gutensteiner Kalken (Anis) des Rabenkogels (Steirisches Salzkammergut, Österreich). *Jahrbuch der geologischen Bundesanstalt*, 143: 605–612.

- Sokač, B., 1968. A new genus of calcareous algae from the Middle Triassic of Velebit. *Geološi Vjesnik*, 21: 207–212.
- Sokač, B. & Velić, I., 1981. *Humiella teutae* n. gen. n. sp. (Dasycladaceae) from the Neocomian of Southern Herzegovina. *Geološi Vjesnik*, 33: 101–105.
- Stachacz, M. & Matysiak, M., 2020. Early Middle Triassic (Anisian) trace fossils, ichnofabrics, and substrate types from the south-eastern Germanic Basin (Wellenkalk facies) of Upper Silesia, southern Poland: Implications for biotic recovery following the Permian/Triassic mass extinction. *Global and Planetary Change*, 194, 103290.
- Steinmann, G., 1903. "Tetraporella remesi", eine neue Dasycladacea aus dem Tithon von Stramberg. *Beiträge zur Paläontologie und Geologie Österreich-Ungarns und des Orients*, 15: 45–54.
- Szulc, J., 1989. Shallow-water carbonate basins of seismically active zones; the example of the Muschelkalk Basin of the Upper Silesia. *Przegląd Geologiczny*, 37: 248–252. [In Polish, with English summary.]
- Szulc, J., 1993. Early Alpine tectonics and lithofacies succession in the Silesian part of the Muschelkalk basin. A Synopsis. In: Hagdorn, H. & Seilacher, A. (eds), *Muschelkalk. Schöntaler Symposium 1991*. Goldschneck-Verlag, Werner K. Weidert, Korb, pp. 19–28.
- Szulc, J., 2000. Middle Triassic evolution of the northern PeriTethys area as influenced by early opening of the Tethys Ocean. *Annales Societatis Geologorum Poloniae*, 70: 1–48.
- Śliwiński, S., 1961. Warstwy olkuskie. *Rudy i Metale Nieżelazne*, 12: 526–529. [In Polish.]
- Wyczółkowski, J., 1971. Effect of the surface morphology of the Palaeozoic substratum on the sedimentation of the Bunter Sandstone and Lower Muschelkalk. *Bulletyn Instytutu Geologicznego*, 243: 121–150. [In Polish, with English summary.]
- Zawidzka, K., 1975. Conodont stratigraphy and sedimentary environment of the Muschelkalk in Upper Silesia. *Acta Geologica Polonica*, 25: 217–256.

Appendix

GPS coordinates of all outcrops presented in this study (in alphabetical order)

- Dąbrowa Górnica – “Ząbkowice” active quarry (50°22'19"N, 19°17'54"E)
- Farska Kolonia – five small inactive quarries (50°32'18"N, 18°16'53"E; 50°32'09"N, 18°16'50"E; 50°32'16"N, 18°16'15"E; 50°31'52"N, 18°16'36"E; 50°32'09"N, 18°16'40"E)
- Jaroszowiec – “Stare Gliny” active quarry (50°21'05"N, 19°35'22"E)
- Kamień Śląski – active quarry (50°33'31"N, 18°07'53"E)
- Libiąż – active quarry (50°06'51"N, 19°20'04"E)
- Nowa Wioska – “GZD” active quarry (50°30'01"N, 19°12'40"E)
- Nowa Wioska – “PROMAG” active quarry (50°30'35"N, 19°14'02"E)
- Nowa Wioska – “TRIBAG” active quarry (50°30'05"N, 19°14'16"E)
- Przelajka – not existing quarry (50°22'33"N, 19°03'09"E)
- Strzelce Opolskie – active quarry (50°31'44"N, 19°18'43"E)
- Tarnów Opolski – active quarry (50°33'11"N, 18°05'34"E)

